

A scientific note on the lactic acid bacterial flora in honeybees in the USA – A comparison with bees from Sweden*

Alejandra VÁSQUEZ¹, Tobias C. OLOFSSON¹, Diana SAMMATARO²

¹ Microbiology Laboratory at Campus Helsingborg, Department of Cell and Organism Biology, Lund University Campus Helsingborg, Rönnowsgatan 6, PO Box SE-25108 Helsingborg, Sweden

² USDA-ARS Carl Hayden Honey Bee Research Centre, Department of Entomology, University of Arizona, 2000 East Allen Road, Tucson, AZ 85719, USA

Received 3 June 2008 – Revised 2 September 2008 – Accepted 2 September 2008

Lactobacillus / *Bifidobacterium* / *Apis mellifera* / 16S rRNA genotyping / coevolution

Recently, it was discovered by Olofsson and Vásquez (2008) that a novel flora composed of lactic acid bacteria (LAB) of the genera *Lactobacillus* and *Bifidobacterium* (Fig. 1), exists in the honey stomach of the honeybee *Apis mellifera*. The ten different flora members varied numerically with the sources of nectar and the presence of other bacterial genera within the honeybee. Closely related phylotypes of some of the lactic acid bacteria have been encountered before in previous work (Scardovi and Trovatielli, 1969; Jeyaprakash et al., 2003; Babendreier et al., 2007) but only in the intestines of honeybees. It appeared that honeybees and the novel LAB flora evolved in mutual dependence on one another, the LAB obtaining a niche in which nutrients were available, the honeybees and their honey in turn being protected by the LAB from harmful microorganisms. In addition, four novel bacterial phylotypes belonging to the family *Pasteurellaceae*, were discovered (Fig. 1). In other studies, (Jeyaprakash et al., 2003; Babendreier et al., 2007) clones with high sequence similarities have been found (Fig. 1), but again only in the intestines of honeybees.

By comparison, honey stomachs of honeybees (*Apis mellifera* and *A. m. scutellata*) collected in the USA were sampled. During spring, honeybees were foraging on flowers that were in bloom (see Tab. I)

around the Carl Hayden Honey Bee Lab (Tucson, AZ) which has over 50 colonies of bees on location. Thirty bees that were actively working flowers were collected from each flower type. Between ten and 20 honey stomachs filled with nectar were removed, placed in an Eppendorf tube with 100 μ L physiological saline kept on ice and refrigerated until express shipped with cold packs to Sweden for bacterial analyzes. In Sweden, a total of 45 cultivated isolates were identified using the 16S rRNA gene sequencing technique (methods of Olofsson and Vásquez, 2008).

The results showed that the Swedish bacterial phylotypes, representing all the groups of *Lactobacillus* and *Bifidobacterium*, were found in the honey stomachs of American bees (Fig. 1). In addition, a novel *Lactobacillus* phylotype 3d was isolated. This phylotype is possibly a novel species belonging to the *Lactobacillus* genus. The 16S rRNA gene sequence of 830 bp resembled the type strain of *Lactobacillus helveticus* DSM 20075^T by 92.3%, which is **below** the threshold level generally used to define a species (95–97%) (Ludwig et al., 1998).

Babendreier et al. (2007) isolated a clone with high sequence similarity to this phylotype but from the intestines of honeybees (Fig. 1). One of the American honey stomach isolates (B4) was identical in sequence to the Swedish *Bifidobacterium* phylotype Trm9 (isolated only from the intestine), which suggests a honey stomach origin.

Olofsson and Vásquez (2008) suggested that the Swedish *Lactobacillus kunkeei* phylotype Fhon2

Corresponding author: T.C. Olofsson,
Tobias.Olofsson@cob.lu.se

* Manuscript editor: Klaus Hartfelder

Table I. Bacterial phylotypes, closely related to the Swedish phylotypes, isolated from American honeybee honey stomachs when bees were foraging on the flowers rosemary (*Rosmarinus officinalis*), wild orange rootstock (*Citrus* spp.), grapefruit (*Citrus paradisi* Red var.) and kumquat (*Fortunella* spp.). Numbers of isolates found, closely related to the Swedish phylotypes by the 16S rRNA gene, are given. Unique American phylotype*.

Phylotype	Rosemary	Bitter orange	Grapefruit	Kumquat
Fhon2	2	2	7	6
Hon2		3	3	
Bma5			1	
Hma2		1	1	
3d*			6	1
Bin7		1	2	
Hma3		1	2	1
Trm9			1	
Bma6		1		1
Lv2			2	

ACKNOWLEDGEMENTS

This study was financed by Gyllenstiernska Krapperupstiftelsen, Ekhagastiftelsen and Sparbankstiftelsen Skåne. We are grateful for the help of the beekeeper Tage Kimblad, late ass. Prof. Sten Ståhl and others for their knowledge and experience, and for their comments and reflections on our work.

Note scientifique sur la flore des bactéries d'acide lactique chez les abeilles domestiques aux États-Unis – Comparaison avec les abeilles de Suède.

Eine wissenschaftliche Notiz zur Milchsäurebakterienflora in Honigbienen aus den USA – Ein Vergleich mit Bienen aus Schweden.

REFERENCES

- Babendreier D., Joller D., Romeis J., Bigler F., Widmer F. (2007) Bacterial community structures in honeybee intestines and their response to two insecticidal proteins, *FEMS Microbiol. Ecol.* 59, 600–610.
- Jeyaprakash A., Hoy M.A., Allsopp M.H. (2003) Bacterial diversity in worker adults of *Apis mellifera capensis* and *Apis mellifera scutellata* (Insecta: Hymenoptera) assessed using 16S rRNA sequences, *J. Invertebr. Pathol.* 84, 96–103.
- Ludwig W., Strunk O., Klugbauer S., Klugbauer N., Weizenegger M., Neumaier J., Bachleitner M., Schleifer K.H. (1998) Bacterial phylogeny based on comparative sequence analysis, *Electrophoresis* 19, 554–568.
- Olofsson T.C., Vásquez A. (2008) Detection and identification of a novel lactic acid bacterial flora within the honey stomach of the honeybee *Apis mellifera*, *Curr. Microbiol.* 57, 356–363.
- Scardovi V., Trovatielli L.D. (1969) New species of bifid bacteria from *Apis mellifica* L. and *Apis indica* F. A contribution to the taxonomy and biochemistry of the genus *Bifidobacterium*, *Zentralbl. Bakteriol. Parasitenkd. Infektionskr. Hyg.* 123, 64–88.